

Totally asymmetric potential

Q: [1] pr 4.11 (a) Given a time reversal invariant Hamiltonian, show that for any energy eigenket

$$\langle \mathbf{L} \rangle = 0. \quad (1.1)$$

(b) If the wave function of such a state is expanded as

$$\sum_{l,m} F_{lm} Y_{lm}(\theta, \phi), \quad (1.2)$$

what are the phase restrictions on F_{lm} ?

A: part (a) For a time reversal invariant Hamiltonian H we have

$$H\Theta = \Theta H. \quad (1.3)$$

If $|\psi\rangle$ is an energy eigenstate with eigenvalue E , we have

$$\begin{aligned} H\Theta |\psi\rangle &= \Theta H |\psi\rangle \\ &= \lambda \Theta |\psi\rangle, \end{aligned} \quad (1.4)$$

so $\Theta |\psi\rangle$ is also an eigenvalue of H , so can only differ from $|\psi\rangle$ by a phase factor. That is

$$\begin{aligned} |\psi'\rangle &= \Theta |\psi\rangle \\ &= e^{i\delta} |\psi\rangle. \end{aligned} \quad (1.5)$$

Now consider the expectation of \mathbf{L} with respect to a time reversed state

$$\begin{aligned} \langle \psi' | \mathbf{L} | \psi' \rangle &= \langle \psi | \Theta^{-1} \mathbf{L} \Theta | \psi \rangle \\ &= \langle \psi | (-\mathbf{L}) | \psi \rangle, \end{aligned} \quad (1.6)$$

however, we also have

$$\begin{aligned} \langle \psi' | \mathbf{L} | \psi' \rangle &= \left(\langle \psi | e^{-i\delta} \right) \mathbf{L} \left(e^{i\delta} | \psi \rangle \right) \\ &= \langle \psi | \mathbf{L} | \psi \rangle, \end{aligned} \quad (1.7)$$

so we have $\langle \psi | \mathbf{L} | \psi \rangle = -\langle \psi | \mathbf{L} | \psi \rangle$ which is only possible if $\langle \mathbf{L} \rangle = \langle \psi | \mathbf{L} | \psi \rangle = 0$.

A: part (b) Consider the expansion of the wave function of a time reversed energy eigenstate

$$\begin{aligned}\langle \mathbf{x} | \Theta | \psi \rangle &= \langle \mathbf{x} | e^{i\delta} | \psi \rangle \\ &= e^{i\delta} \langle \mathbf{x} | \psi \rangle ,\end{aligned}\tag{1.8}$$

and then consider the same state expanded in the position basis

$$\begin{aligned}\langle \mathbf{x} | \Theta | \psi \rangle &= \langle \mathbf{x} | \Theta \int d^3 \mathbf{x}' (|\mathbf{x}'\rangle \langle \mathbf{x}'|) | \psi \rangle \\ &= \langle \mathbf{x} | \Theta \int d^3 \mathbf{x}' (\langle \mathbf{x}' | \psi \rangle) |\mathbf{x}'\rangle \\ &= \langle \mathbf{x} | \int d^3 \mathbf{x}' (\langle \mathbf{x}' | \psi \rangle)^* \Theta |\mathbf{x}'\rangle \\ &= \langle \mathbf{x} | \int d^3 \mathbf{x}' (\langle \mathbf{x}' | \psi \rangle)^* |\mathbf{x}'\rangle \\ &= \int d^3 \mathbf{x}' (\langle \mathbf{x}' | \psi \rangle)^* \langle \mathbf{x} | \mathbf{x}' \rangle \\ &= \int d^3 \mathbf{x}' \langle \psi | \mathbf{x}' \rangle \delta(\mathbf{x} - \mathbf{x}') \\ &= \langle \psi | \mathbf{x} \rangle .\end{aligned}\tag{1.9}$$

This demonstrates a relationship between the wave function and its complex conjugate

$$\langle \mathbf{x} | \psi \rangle = e^{-i\delta} \langle \psi | \mathbf{x} \rangle .\tag{1.10}$$

Now expand the wave function in the spherical harmonic basis

$$\begin{aligned}\langle \mathbf{x} | \psi \rangle &= \int d\Omega \langle \mathbf{x} | \hat{\mathbf{n}} \rangle \langle \hat{\mathbf{n}} | \psi \rangle \\ &= \sum_{lm} F_{lm}(r) Y_{lm}(\theta, \phi) \\ &= e^{-i\delta} \left(\sum_{lm} F_{lm}(r) Y_{lm}(\theta, \phi) \right)^* \\ &= e^{-i\delta} \sum_{lm} (F_{lm}(r))^* Y_{lm}^*(\theta, \phi) \\ &= e^{-i\delta} \sum_{lm} (F_{lm}(r))^* (-1)^m Y_{l,-m}(\theta, \phi) \\ &= e^{-i\delta} \sum_{lm} (F_{l,-m}(r))^* (-1)^m Y_{l,m}(\theta, \phi),\end{aligned}\tag{1.11}$$

so the F_{lm} functions are constrained by

$$F_{lm}(r) = e^{-i\delta} (F_{l,-m}(r))^* (-1)^m .\tag{1.12}$$

Bibliography

- [1] Jun John Sakurai and Jim J Napolitano. *Modern quantum mechanics*. Pearson Higher Ed, 2014. 1